

Sparse modelling and multi-exponential analysis

Dagstuhl Seminar 15251

Organizers:

Annie Cuyt
University of Antwerp, Belgium

George Labahn
University of Waterloo, Canada

Avraham Sidi
Technion – Haifa, Israel

Coordinator:

Wen-shin Lee
University of Antwerp, Belgium

June 14–19 , 2015



SCHLOSS DAGSTUHL
Leibniz-Zentrum für Informatik

Contents

Seminar Program	7
Accuracy of Spike-Train Fourier Reconstruction for Near-Colliding Nodes	13
<i>Andrey Akinshin, Yosef Yomdin, Dmitry Batenkov</i>	
Multidimensional approximation of functions sampled at unequally spaced points by sums of exponentials	14
<i>Fredrik Andersson, Marcus Carlsson, Herwig Wendt</i>	
Fourier-Sparsity Testing of Boolean Functions	16
<i>Andrew Arnold, Eric Blais</i>	
Error-Correcting Sparse Interpolation in Chebyshev Basis	17
<i>Andrew Arnold, Erich L. Kaltofen</i>	
Numerical stability of the parameter estimation problem in sparse generalized exponential sums	18
<i>Dmitry Batenkov</i>	
On the conditioning of the Padé map and related questions	19
<i>Bernhard Beckermann</i>	
Sub-Nyquist spectral analysis	20
<i>Matteo Briani, Annie Cuyt, Wen-shin Lee</i>	
Order parameter for images of structured arrays	21
<i>Adhemar Bultheel, Forrest Kaatz</i>	
A moment matrix approach to symmetric cubatures	22
<i>Mathieu Collowald, Evelyne Hubert</i>	
A Sparse Sampling Method to Estimate Parameters in Multivariate Exponential Sums	23
<i>Annie Cuyt, Wen-shin Lee</i>	
Exponential analysis, Sparse interpolation and Padé approximation	24
<i>Annie Cuyt, Wen-shin Lee</i>	
Trivariate polynomial approximation on Lissajous curves	25
<i>Stefano De Marchi</i>	
Inverse Problems regularised by Sparsity	26
<i>Pier Luigi Dragotti</i>	
Mobile Sampling	27
<i>Karlheinz Gröchenig</i>	
High dimensional approximation with trigonometric polynomials	28
<i>Lutz Kämmerer, Daniel Potts, Toni Volkmer</i>	

A multivariate generalization of Prony’s method	29
<i>Stefan Kunis, Thomas Peter, Tim Römer, Ulrich von der Ohe</i>	
Behavior preserving extension of univariate and bivariate functions	30
<i>David Levin</i>	
High-speed fluorescence lifetime imaging (FLIM) instruments	31
<i>David Li</i>	
Estimating Variable Star Periods from Unevenly Sampled Light Curve Data	32
<i>Daniel Lichtblau</i>	
Structured low-rank approximation: Theory, algorithms, and applications	33
<i>Ivan Markovsky</i>	
Well conditioned rational functions approximants versus numerically coprime polynomials	34
<i>Ana C. Matos</i>	
Using noise to detect faint signals: tricks with Padé approximants to Z-transforms	35
<i>Luca Perotti, Daniel Bessis, Daniel Vrinceanu, Tania Regimbau</i>	
The generalized Prony method and its application I and II	36
<i>Thomas Peter, Gerlind Plonka</i>	
Efficient spectral estimations by MUSIC and related algorithms	37
<i>Daniel Potts, Toni Volkmer, Manfred Tasche</i>	
Using univariate algorithms to solve multivariate problems	38
<i>Daniel S. Roche</i>	
New approaches to Vector-Valued Rational Interpolation	39
<i>Avram Sidi</i>	
Towards simplified construction of subresultant matrix of multiple univariate polynomials	41
<i>Akira Terui</i>	
Hankel and Quasi-Hankel low-rank matrix completion: a convex relaxation	42
<i>Konstantin Usevich, Pierre Comon</i>	
A deterministic sparse FFT algorithm for vectors with short support	43
<i>Katrin Wannowwetsch</i>	
Sparsity with Symbolic Polynomials	44
<i>Stephen M. Watt</i>	

Reconstruction of Structured Functions from Sparse Fourier Data	45
<i>Marius Wischerhoff</i>	
Semidefinite Representations of Noncompact Convex Sets	46
<i>Lihong Zhi</i>	
List of Participants	47

Seminar Program

Monday June 15, 2015

Opening

- 10.00–10.30 Opening
- 10.30–11.00 **Exponential analysis, Sparse interpolation and Padé approximation**
Annie Cuyt, Wen-shin Lee
- 11.00–12.00 Discussion
- 12.15–13.45 Lunch

Application Section

- 14.30–15.00 **Mobile Sampling**
Karlheinz Gröchenig
- 15.00–15.30 **Order parameter for images of structured arrays**
Adhemar Bultheel, Forrest Kaatz
- 15.30–16.00 **Sub-Nyquist spectral analysis**
Matteo Briani, Annie Cuyt, Wen-shin Lee
- 16.00–16.30 Coffee break
- 16.30–17.00 **High-speed fluorescence lifetime imaging (FLIM) instruments**
David Li
- 17.00–17.30 **Inverse Problems regularised by Sparsity**
Pier Luigi Dragotti
- 17.30–18.00 **Estimating Variable Star Periods from Unevenly Sampled Light Curve Data**
Daniel Lichtblau
- 18.00–19.30 Dinner

Tuesday June 16, 2015

Exponential Analysis

- 08.45–09.45 Discussion
- 09.45–10.15 **The generalized Prony method and its application I and II**
Thomas Peter, Gerlind Plonka
- 10.15–10.45 **The generalized Prony method and its application I and II**
Thomas Peter, Gerlind Plonka
- 10.45–11.15 Coffee break
- 11.15–11.45 **A multivariate generalization of Prony’s method**
Stefan Kunis, Thomas Peter, Tim Römer, Ulrich von der Ohe
- 11.45–12.15 **A multivariate generalization of Prony’s method**
Stefan Kunis, Thomas Peter, Tim Römer, Ulrich von der Ohe
- 12.15–13.45 Lunch

Exponential Analysis

- 13.45–14.30 Discussion
- 14.30–15.00 **Numerical stability of the parameter estimation problem in sparse generalized exponential sums**
Dmitry Batenkov
- 15.00–15.30 **A deterministic sparse FFT algorithm for vectors with short support**
Katrin Wannewetsch
- 15.30–16.00 **Reconstruction of Structured Functions from Sparse Fourier Data**
Marius Wischerhoff
- 16.00–16.30 Coffee break
- 16.30–17.00 **Accuracy of Spike-Train Fourier Reconstruction for Near-Colliding Nodes**
Andrey Akinshin, Yosef Yomdin, Dmitry Batenkov
- 17.00–17.30 **High dimensional approximation with trigonometric polynomials**
Lutz Kämmerer, Daniel Potts, Toni Volkmer
- 17.30–18.00 **Efficient spectral estimations by MUSIC and related algorithms**
Daniel Potts, Toni Volkmer, Manfred Tasche
- 18.00–19.30 Dinner

Wednesday June 17, 2015

Approximation Theory

- 09.00–09.30 **New approaches to Vector-Valued Rational Interpolation**
Avram Sidi
- 09.30–10.00 **Using noise to detect faint signals: tricks with Padé approximants to \mathbf{Z} -transforms**
Luca Perotti, Daniel Bessis, Daniel Vrinceanu, Tania Regimbau
- 10.00–10.30 **Trivariate polynomial approximation on Lissajous curves**
Stefano De Marchi
- 10.30–11.00 Coffee break
- 11.00–11.30 **Behavior preserving extension of univariate and bivariate functions**
David Levin
- 11.30–12.00 **On the conditioning of the Padé map and related questions**
Bernhard Beckermann
- 12.00–12.30 **Well conditioned rational functions approximants versus numerically co-prime polynomials**
Ana C. Matos
- 12.30–13.45 Lunch
- Excursion
- 18.00–19.30 Dinner

Thursday June 18, 2015

Sparse Interpolation

08.45–09.45 Discussion

09.45–10.15 **Error-Correcting Sparse Interpolation in Chebyshev Basis**

Andrew Arnold, Erich L. Kaltofen

10.15–10.45 **Multidimensional approximation of functions sampled at unequally spaced points by sums of exponentials**

Fredrik Andersson, Marcus Carlsson, Herwig Wendt

10.45–11.15 Coffee break

11.15–11.45 **Using univariate algorithms to solve multivariate problems**

Daniel S. Roche

11.45–12.15 **A Sparse Sampling Method to Estimate Parameters in Multivariate Exponential Sums**

Annie Cuyt, Wen-shin Lee

12.15–13.45 Lunch

Sparse Interpolation

13.45–14.30 Discussion

14.30–15.00 **Sparsity with Symbolic Polynomials**

Stephen M. Watt

15.00–15.30 **Fourier-Sparsity Testing of Boolean Functions**

Andrew Arnold, Eric Blais

15.30–16.00 **Towards simplified construction of subresultant matrix of multiple univariate polynomials**

Akira Terui

16.00–16.30 Coffee break

16.30–18.00 Discussion

18.00–19.30 Dinner

Friday June 19, 2015

Related Topics

08.45–09.45 Discussion

09.45–10.15 **Structured low-rank approximation: Theory, algorithms, and applications**
Ivan Markovsky

10.15–10.45 **Hankel and Quasi-Hankel low-rank matrix completion: a convex relaxation**
Konstantin Usevich, Pierre Comon

10.45–11.15 Coffee break

11.15–11.45 **Semidefinite Representations of Noncompact Convex Sets**
Lihong Zhi

11.45–12.15 **A moment matrix approach to symmetric cubatures**
Mathieu Collowald, Evelyne Hubert

12.15–13.45 Lunch

Departure

Accuracy of Spike-Train Fourier Reconstruction for Near-Colliding Nodes

Andrey Akinshin Yosef Yomdin
The Weizmann Institute of Science
Rehovot
Israel

andrey.akinshin@gmail.com, yosef.yomdin@weizmann.ac.il

Dmitry Batenkov
Technion - Israel Institute of Technology
Haifa 32000
Israel
batenkov@cs.technion.ac.il

Abstract

We study reconstruction of “spike-train” signals F of the form $F(x) = \sum_{j=1}^d a_j \delta(x-x_j)$, from their Fourier transform $\hat{F}(s)$, known for $s \in [-N, N]$, with an absolute error not exceeding $\epsilon > 0$. We concentrate on “near-collision” situations where the nodes x_j are known to form an l elements cluster of a size $h \ll 1$.

We show that in such situations the geometry of error amplification in the reconstruction is governed by the “Prony foliations” S_q whose leaves are defined by the Prony equations $\sum_{j=1}^d a_j x_j^k = \gamma_k$, with $k = 0, \dots, q \leq l$, and with the arbitrary right-hand sides γ_k . On this base we give an “absolute” (i.e. valid with any reconstruction method) lower bound for the “worst case” reconstruction error of F from \hat{F} . We show that for the measurement error $\epsilon > C_1(hN)^{2l-1}$, the inside configuration of the cluster nodes (in the worst case scenario) cannot be reconstructed at all.

Combining a proper rescaling with the “Decimation method” we show that for $\epsilon < C_2(hN)^{2l-1}$, $C_2 \ll C_1$, an accurate (up to an error αh , $\alpha \ll 1$) reconstruction of the cluster nodes is possible. The same algorithm reconstructs the non-cluster nodes and amplitudes with the full accuracy (of order $\frac{\epsilon}{N}$).

Multidimensional approximation of functions sampled at unequally spaced points by sums of exponentials

Fredrik Andersson Marcus Carlsson
 Centre for Mathematical Sciences
 Lund University
 Sweden
 {fa,mc}@maths.lth.se

Herwig Wendt
 IRIT-ENSEEITH
 University of Toulouse
 France
 herwig.wendt@irit.fr

Abstract

Let f be sampled at unequally spaced points $x_m \in \mathbb{R}^d$. We consider the problem of finding

$$g(x) = \sum_{k=1}^K c_k e^{2\pi i x \cdot \xi_k}, \quad (1)$$

so that $f(x_m) \approx g(x_m)$.

Let Ξ and Υ be subsets of equally spaced grids in \mathbb{R}^d and let $\Omega = \Xi + \Upsilon = \{x + y : x \in \Xi, y \in \Upsilon\}$. Given a function g on Ω , consider the generalized multidimensional Hankel operator

$$\Gamma_g h(x) = \sum_{y \in \Upsilon} g(x + y) h(y), \quad x \in \Xi \quad (2)$$

By Kronecker's theorem Γ_g has rank K if g is of the form (1). It also turns out that range of Γ_g is the space of all linear combinations of the functions $e^{2\pi i x \cdot \xi_k}$ on Ξ (See Lemma 4.2 of [1]). Let us represent the operator with the matrix $\mathbf{\Gamma}_g$.

Let \mathbf{J} be an interpolation matrix that interpolates the values at the equally spaced point in Ω to unequally spaced points $\Psi = \{x_m\}_{m=1}^M$ in \mathbb{R}^d . In order to approximate the function f sampled at Ψ using K exponentials, we consider the optimization problem

$$\begin{aligned} & \underset{g}{\text{minimize}} && \sum_{m=1}^M |(\mathbf{J}g)_m - f(x_m)|^2 \\ & \text{subject to} && \text{rank } \mathbf{\Gamma}_g = K \end{aligned} \quad (3)$$

where $(\mathbf{J}g)_m$ is the interpolated value of g at x_m .

We follow the setup in [4], and formulate (3) using the alternating direction method of multipliers [5]. The problem formulations is not convex, and there is hence no guarantee that the procedure will converge. However, it will typically give a matrix values of g such that the singular values σ_k of $\mathbf{\Gamma}_g$ are small if $k > K$. To estimate the (multidimensional) frequencies ξ_k associated with g we can then follow the approach gives in [2, 3] by solving systems of polynomial equations with coefficients taken from the singular vectors of $\mathbf{\Gamma}_g$ for $k > K$.

References

- [1] Fredrik Andersson and Marcus Carlsson. On general domain truncated correlation and convolution operators with finite rank. *Integral Equations and Operator Theory*, pages 1-32, 2015.
- [2] Fredrik Andersson, Marcus Carlsson, and V Maarten. Nonlinear approximation of functions in two dimensions by sums of exponential functions. *Applied and Computational Harmonic Analysis*, 29(2):156-181, 2010.
- [3] Fredrik Andersson, Marcus Carlsson, and V Maarten. Nonlinear approximation of functions in two dimensions by sums of wave packets. *Applied and Computational Harmonic Analysis*, 29(2):198-213, 2010.
- [4] Fredrik Andersson, Marcus Carlsson, Jean-Yves Tournet, and Herwig Wendt. A new frequency estimation method for equally and unequally spaced data. *Signal Processing, IEEE Transactions on*, 62(21):5761-5774, 2014.
- [5] Stephen Boyd, Neal Parikh, Eric Chu, Borja Peleato, and Jonathan Eckstein. Distributed optimization and statistical learning via the alternating direction method of multipliers. *Foundation and Trends in Machine Learning*, 3(1):1-122, 2011.

Fourier-Sparsity Testing of Boolean Functions

Andrew Arnold Eric Blais
University of Waterloo
Waterloo, Ontario
Canada
{a4arnold,eblais}@uwaterloo.ca

Abstract

We consider the problem of testing whether a function f has at most s nonzero Fourier coefficients, in which case we say f is s -sparse, given black-box access to f . We restrict our attention to perhaps the simplest case when f is a Boolean function acting on n bits. The analogous problem of learning the Fourier transform of an s -sparse Boolean function f was studied in previous work by Kushilevitz and Mansour [SIAM J. Computing, vol 22. (1992)], and Levin [J. Symb. Logic, vol 58. (1993)], the latter resulting in an $O(ns)$ Monte Carlo Sparse Fourier Transform (SFT) algorithm. Their work was the foundation for subsequent Sparse Fourier Transform algorithms in more general settings.

We say an algorithm is an ϵ -tester for sparse Boolean functions if it accepts if f is s -sparse and rejects if f is ϵ -far from s -sparse in terms of ℓ_2 norm, each with probability at least $2/3$. Gopalan et al. [SIAM J. Computing, vol 40. (2011)], gave the first such tester with query-complexity polynomial in s and ϵ^{-1} .

We improve upon this result, present a sparsity tester with query-complexity $O(s \log s \epsilon^{-2} + \epsilon^{-4})$. Our tester relies on dimensionality-reduction techniques developed in the aforementioned previous work. Using these techniques, we reduce sparsity testing to the problem of homomorphism testing, which in turn may be solved via the Blum-Luby-Rubinfeld (BLR) linearity test [J. Comput. Syst. Sci. Int., vol 47. (1993)].

Error-Correcting Sparse Interpolation in Chebyshev Basis

Andrew Arnold*
University of Waterloo
Waterloo, Ontario
Canada
a4arnold@uwaterloo.ca

Erich L. Kaltofen†
North Carolina State University
Raleigh, North Carolina
USA
kaltofen@ncsu.edu

Abstract

We present an error-correcting interpolation algorithm for a univariate black-box polynomial that has a sparse representation using Chebyshev polynomials as a term basis. Our algorithm assumes that an upper bound on the number of erroneous evaluations is given as input, and is a generalization of the algorithm by Lakshman and Saunders [SIAM J. Comput., vol. 24 (1995)] for interpolating sparse Chebyshev polynomials and the techniques in error-correcting sparse interpolation in the usual basis of consecutive powers of the variable due to Comer, Kaltofen, and Pernet [Proc. ISSAC 2012 and 2014]. We prove the correctness of our list-decoder-based algorithm with a Descartes-rule-of-signs-like property for sparse polynomials in Chebyshev basis. We also give a new algorithm that reduces the sparse interpolation in Chebyshev basis to that in power basis, thus making the many techniques for the sparse interpolation in power basis, for instance, supersparse (lacunary) interpolation over large finite fields, available to interpolation in Chebyshev basis. Furthermore, we can customize the randomized early termination algorithms from Kaltofen and Lee [J. Symb. Comput., vol. 36 (2003)] to our new approach.

*This research was supported in part by NSERC.

†This research was supported in part by the National Science Foundation under Grants CCF-1115772 and CCF-1421128

Numerical stability of the parameter estimation problem in sparse generalized exponential sums

Dmitry Batenkov*

Technion – Israel Institute of Technology

Haifa 32000

Israel

batenkov@cs.technion.ac.il

Abstract

We consider the parameter estimation problem in sparse generalized exponential sums of the form $m(k) = \sum_{j=1}^s e^{ix_j k} \sum_{\ell=0}^{d_j-1} a_{\ell,j} k^\ell$, when $m(k)$ are known only approximately.

We provide estimates on the component-wise condition numbers of the parameters x_j and $a_{\ell,j}$ above, and show that they can be accurately recovered by sampling at arithmetic progressions and polynomial homotopy methods.

We also discuss the application of these ideas to the problem of recovering a piecewise-smooth function (including the positions of the discontinuities) from its Fourier coefficients.

References

- [1] D. Batenkov. Complete Algebraic Reconstruction of Piecewise-Smooth Functions from Fourier Data. *Mathematics of Computation*, 2015.
- [2] Dmitry Batenkov. Accurate solution of near-colliding Prony systems via decimation and homotopy continuation. *arXiv:1501.00160 [cs, math]*, December 2014. arXiv: 1501.00160.

*The research leading to these results has received funding from the European Research Council under European Union's Seventh Framework Program, ERC Grant agreement no. 320649

On the conditioning of the Padé map and related questions

Bernhard Beckermann*

Laboratoire Painlevé UMR 8524, UFR Mathématiques

Université de Lille, France

{bbecker,matos}@math.univ-lille1.fr

Abstract

Padé approximants play an important role in signal processing, sparse interpolation and exponential analysis. In this talk we will report about recent results from [1] concerning the forward and backward conditioning of the (real) Padé map, which sends a vector of Taylor coefficients onto the normalized vector of coefficients of the Padé numerator and denominator. In particular, we show that this map is not necessarily well conditioned for robust Padé approximants in the sense of Trefethen et al [2].

We will also discuss the condition number of related non-linear maps.

This is a joint work with Ana C. Matos.

References

- [1] B. Beckermann, A. Matos, Algebraic properties of robust Padé approximants *Journal of Approx. Theory* **190**, 91-115 (2015)
- [2] P. Gonnet, S. Güttel and L.N. Trefethen, Robust Padé approximation via SVD, *SIAM Review*, 55 (2013), 101-117.

*Supported in part by the Labex CEMPI (ANR-11-LABX-0007-01).

Sub-Nyquist spectral analysis

Matteo Briani Annie Cuyt Wen-shin Lee
Dept. Mathematics and Computer Science
University of Antwerp
Belgium

{matteo.briani, annie.cuyt, wen-shin.lee}@uantwerpen.be

Abstract

In the field of sparse interpolation, parametric methods aim to retrieve the values of parameters of a linear combination of exponential functions from samples in a uniform time grid. These samples are collected following the Shannon-Nyquist theorem that dictates the minimum sampling rate that prevents the aliasing effect. In this paper we explain how it is possible, by means of undersampling, to use a coarser time grid and still be able to solve the aliasing effect. This reflects into a better conditioning of the problem and this behavior is explained by means of the ill-disposedness and a link to Padé approximation theory. Avoiding the aliasing effect, and using a coarser time grid, it is possible to perform several smaller independent analysis from the original set of samples. Joining these analysis together we obtain a method that brings higher accuracy to the existing parametric methods and introduces an extra parameter that can be use as validator. This is especially useful when the parametric method has to deal with signals consisting of close frequencies in a broad spectrum.

Order parameter for images of structured arrays

Adhemar Bultheel
Dept. Computer Science, KU Leuven
Heverlee
Belgium
adhemar.bultheel@kuleuven.be

Forrest Kaatz
Mesalands Community College
Tucumcari
NM, USA
fhkaatz@gmail.com

Abstract

In nature (e.g. a bee honeycomb, muscle structure, crystals) or in engineering (e.g. micro lens arrays, nano pore/pillar arrays, solar cells) two-dimensional highly regular arrays are produced. Hexagonal, square or triangular grids are most common. Perfect symmetry of the grids does not exist in practical situations. Given the image of some array, one may analyse the properties of each of the individual nodes of the grid and compute parameters like their size, the location of their centers, perhaps their orientation, etc. These parameters could be combined to define some number indicating the deviation from the ideal grid.

We have tried to compute some order parameter from the Fourier transform of the image. For example a perfect hexagonal array has a Fourier spectrum that consists of a central peak, surrounded by six smaller peaks and their harmonics. This is a sparse exponential representation. The more the nodes in the image are dislocated from the perfect grid, the more noise will show up in the spectrum. Thus the amount of noise in the Fourier domain can be used as a measure for the disorder of the original grid.

Unfortunately, images may depend on many parameters (number of nodes, size of the nodes, shape of the nodes, orientation of the image, . . .) so that the Fourier technique only works in a rather restrictive number of situations and it is probably not useful in practical situations.

A moment matrix approach to symmetric cubatures

Mathieu Collowald Evelyne Hubert
Inria Méditerranée
Sophia Antipolis, France
{mathieu.collowald, evelyne.hubert}@inria.fr

Abstract

Quadrature and sparse interpolation are closely linked. The common key issue is the construction of a linear form

$$\Lambda : \mathbb{R}[x] \rightarrow \mathbb{R}, p \mapsto \sum_{j=1}^r a_j p(\xi_j)$$

from the knowledge of its restriction to $\mathbb{R}[x]_{\leq d}$. The unknowns are the weights a_j and the nodes ξ_j .

Cubature is a generalization of quadrature in higher dimension. An approach based on moment matrices was proposed in [2, 4]. We give a basis-free version in terms of the Hankel operator \mathcal{H} associated to Λ . The existence of a cubature of degree d with r nodes boils down to conditions of ranks and positive semidefiniteness on \mathcal{H} . The nodes are then the solutions of a generalized eigenvalue problem.

Standard domains of integration are symmetric under the action of a finite group. It is natural to look for cubatures that respect this symmetry [1, 3]. Introducing adapted bases obtained from representation theory, the symmetry constraint allows to block diagonalize the Hankel operator \mathcal{H} . The size of the blocks is explicitly related to the orbit types of the nodes. From the computational point of view, we then deal with smaller-sized matrices both for securing the existence of the cubature and computing the nodes.

References

- [1] R. Cools. Constructing cubature formulae: the science behind the art. *Acta numerica*, 6:1–54, 1997.
- [2] L. Fialkow and S. Petrovic. A moment matrix approach to multivariable cubature. *Integral Equations Operator Theory*, 52(1):85–124, 2005.
- [3] K. Gatermann. The construction of symmetric cubature formulas for the square and the triangle. *Computing*, 40(3):229–240, 1988.
- [4] J. B. Lasserre. The existence of Gaussian cubature formulas. *J. Approx. Theory*, 164(5):572–585, 2012.

A Sparse Sampling Method to Estimate Parameters in Multivariate Exponential Sums

Annie Cuyt Wen-shin Lee
Dept. Mathematics and Computer Science
University of Antwerp
Belgium
{annie.cuyt, wen-shin.lee}@uantwerpen.be

Abstract

We consider the interpolation of an n -variate exponential sum

$$F(x_1, \dots, x_n) = \sum_{j=1}^t c_j e^{f_{j,1}x_1 + f_{j,2}x_2 + \dots + f_{j,n}x_n}.$$

That is, to recover parameters $c_j, f_{j,k}$ from the evaluations of $F(x_1, \dots, x_n)$.

We present a parametric method that can interpolate $F(x_1, \dots, x_n)$ from $(n+1) \cdot t$ evaluations. In order to solve for the multivariate parameters, the target function $F(x_1, \dots, x_n)$ is evaluated at the additional points defined by the *identification shifts*. Since the total number of parameters c_j and $f_{j,k}$ is exactly $(n+1) \cdot t$, our method can interpolate $F(x_1, \dots, x_n)$ from the minimum possible number of evaluations.

In general, our method can be used in exponential analysis problems where additional evaluations of the same exponentials are required. One of such applications is to recover the correct frequencies from the aliased results caused by different subsamplings.

In computer algebra, our idea can be applied to interpolate a blackbox polynomial evaluated at small finite fields.

Our method can be embedded in any Prony-like algorithm, such as the least square Prony, ESPRIT, matrix pencil, etc., thus can be viewed as a new class of tools exposing an additional range of possibilities in exponential analysis.

Exponential analysis, Sparse interpolation and Padé approximation

Annie Cuyt Wen-shin Lee
Dept. Mathematics and Computer Science
University of Antwerp
Belgium
{annie.cuyt, wen-shin.lee}@uantwerpen.be

Abstract

A common underlying problem statement in many applications is that of determining the number of components, and for each component the value of the frequency, damping factor, amplitude and phase in a multi-exponential model. It occurs, for instance, in magnetic resonance and infrared spectroscopy, vibration analysis, seismic data analysis, electronic odour recognition, keystroke recognition, nuclear science, music signal processing, transient detection, motor fault diagnosis, electrophysiology, drug clearance monitoring and glucose tolerance testing, to name just a few.

The general technique of multi-exponential modeling is closely related to what is commonly known as the Padé-Laplace method in approximation theory, and the technique of sparse interpolation in the field of computer algebra. The problem of multi-exponential modeling is an inverse problem and therefore may be severely ill-posed, depending on the relative location of the frequencies and phases. Besides the reliability of the estimated parameters, the sparsity of the multi-exponential representation has become important. A representation is called sparse if it is a combination of only a few elements instead of all available generating elements.

Despite the close connections between these fields, there is a clear lack of communication in the scientific literature. The aim of this seminar is to bring researchers together from the three mentioned fields, with scientists from the varied application domains.

Trivariate polynomial approximation on Lissajous curves

Stefano De Marchi

Department of Mathematics, University of Padova

Italy

demarchi@math.unipd.it

Abstract

We study Lissajous curves in the 3-cube, that generate algebraic cubature formulas on a special family of rank-1 Chebyshev lattices. These formulas are used to construct trivariate hyperinterpolation polynomials via a single 1-d Fast Chebyshev Transform (by the Chebfun package), and to compute discrete extremal sets of Fekete and Leja type for trivariate polynomial interpolation. Applications could arise in the framework of Lissajous sampling for MPI (Magnetic Particle Imaging).

Joint work with Len Bos (University of Verona) and Marco Vianallo (University of Padova)

Inverse Problems regularised by Sparsity

Pier Luigi Dragotti*
Imperial College London
London
United Kingdom
p.dragotti@imperial.ac.uk

Abstract

Modelling signals as sparse in a proper domain has proved useful in many signal processing tasks and, here, we show how sparsity can be used to solve inverse problems. We first recall that many inverse problems involve the reconstruction of continuous-time or continuous-space signals from discrete measurements and show how to relate the discrete measurements to some properties of the original signal (e.g., its Fourier transform at specific frequencies). Given this partial knowledge of the original signal, we then solve the inverse problem using sparsity. We focus on two specific problems which have important practical implications: localisation of diffusion sources from sensor measurements and reconstruction of planar domains from samples. First, we show how to reconstruct specific planar domains whose contours are determined using implicit functions, then we localise diffusion sources using a variation of the ‘reciprocity gap’ method which involves analytic test functions.

In both cases, the problem is solved by building a Prony’s type system and by building structured matrices which, in the ideal settings, are simultaneously Toeplitz and rank deficient.

*Sponsored by European Research Council ERC, project 277800 (RecoSamp). Joint work with: T. Blu (CUHK), M. Vetterli (EPFL), John Murray-Bruce (ICL) and Jon Onativia (ICL)

Mobile Sampling

Karlheinz Gröchenig
Faculty of Mathematics
University of Vienna
Nordbergstrasse 15
A-1090 Vienna, Austria
karlheinz.groechenig@univie.ac.at

Abstract

We study the design of sampling trajectories for stable sampling and the reconstruction of bandlimited spatial fields using mobile sensors. The spectrum is assumed to be a symmetric convex set. As a performance metric we use the path density of the set of sampling trajectories that is defined as the total distance traveled by the moving sensors per unit spatial volume of the spatial region being monitored. Focussing first on parallel lines, we identify the set of parallel lines with minimal path density that contains a set of stable sampling for fields bandlimited to a known set. We then show that the problem becomes ill-posed when the optimization is performed over all trajectories by demonstrating a feasible trajectory set with arbitrarily low path density. However, the problem becomes well-posed if we explicitly specify the stability margins. We demonstrate this by obtaining a non-trivial lower bound on the path density of an arbitrary set of trajectories that contain a sampling set with explicitly specified stability bounds.

This is joint work with Jose Luis Romero, Univ. of Vienna, Jayakrishnan Unnikrishnan and Martin Vetterli from Ecole Polytechnique Fédérale de Lausanne (EPFL), Switzerland.

High dimensional approximation with trigonometric polynomials

Lutz Kämmerer Daniel Potts Toni Volkmer*

Faculty of Mathematics

Technische Universität Chemnitz

D-09107 Chemnitz, Germany

potts@mathematik.tu-chemnitz.de

Abstract

In this talk, we present algorithms for the approximation of multivariate functions by trigonometric polynomials. The approximation is based on sampling of multivariate functions on rank-1 lattices. To this end, we study the approximation of functions in periodic Sobolev spaces of dominating mixed smoothness. The proposed algorithm based mainly on a one-dimensional fast Fourier transform, and the arithmetic complexity of the algorithm depends only on the cardinality of the support of the trigonometric polynomial in the frequency domain. Therefore, we investigate trigonometric polynomials with frequencies supported on hyperbolic crosses and energy based hyperbolic crosses in more detail. Furthermore, we present algorithms where the support of the trigonometric polynomial is unknown.

References

- [1] D. Potts, and T. Volkmer. Sparse high-dimensional FFT based on rank-1 lattice sampling. *Preprint* 2015.
- [2] L. Kämmerer, D. Potts, and T. Volkmer. Approximation of multivariate functions by trigonometric polynomials based on rank-1 lattice sampling. *J. Complexity* 31, 543-576, 2015.

*Sponsored by German Research Foundation within the project PO 711/10-2.

A multivariate generalization of Prony's method

Stefan Kunis Thomas Peter Tim Römer Ulrich von der Ohe*

Institute of Mathematics

Osnabrück University

Germany

{skunis, petert, troemer, uvonderohe}@uos.de

Abstract

A classical solution to the problem of parameter reconstruction for an exponential sum from a finite number of samples is given by Prony's method and the parameters are recovered as roots of a single univariate polynomial. We present a generalization of this method for exponential sums in an arbitrary finite number of variables and realize the parameters as common roots of several multivariate polynomials. Finally, the coefficients of the exponential sum arise as solutions to a linear system of equations.

In the first part of the talk we explain this approach and its algebraic properties. Provided we sample the exponential sum on an equidistant grid with a number of grid points in each coordinate direction bounded from below by the number parameters, unique reconstruction is guaranteed and this bound is shown to be sharp. In its simplest form, the reconstruction method consists of setting up a certain multilevel Toeplitz matrix of the samples, compute a basis of its kernel, and compute by some method of choice the set of common roots of the multivariate polynomials whose coefficients are given in the second step.

The second part of the talk is dedicated to numerical properties of our approach. Provided the number of grid points in each coordinate direction is bounded from below by some small constant divided by the separation distance of the parameters, the kernel of the above Toeplitz matrix can be stably computed. Moreover, we relate our approach to a recent semidefinite optimization formulation and show a couple of numerical experiments.

*Support by DFG-GRK1916 is gratefully acknowledged.

Behavior preserving extension of univariate and bivariate functions

David Levin
Tel Aviv University
Tel Aviv
Israel
levin@tau.ac.il

Abstract

Given function values on a domain D_0 , possibly with noise, we examine the possibility of extending the function to a larger domain D , $D_0 \subset D$. In addition to smoothness at the boundary of D_0 , the extension on $D \setminus D_0$ should also inherit behavioral trends of the function on D_0 , such as growth and decay or even oscillations. The approach chosen here is based upon the framework of linear models, univariate or bivariate, with constant or varying coefficients.

High-speed fluorescence lifetime imaging (FLIM) instruments

David Li

Centre for Biophotonics, University of Strathclyde

Glasgow G4 0RE

UK

David.Li@strath.ac.uk

Abstract

Fast fluorescence lifetime imaging (FLIM) techniques are powerful tools for visualising protein interaction networks in living cells. FLIM has been used for cancer diagnosis, assessing drug efficacy in cancer therapy, understanding brain functions, etc. It can also sense physiological parameters such as Ca^{2+} , pH, O_2 , temperature, viscosity, etc. For real-time applications, such as visualising neuronal activities or fast biophysical phenomena, it is desirable to apply innovative solid-state single-photon sensors and fast hardware embedded exponential analysis processors that can boost FLIM imaging. But is it easy to have a hardware-friendly and high-efficient exponential analysis method for such applications?

Estimating Variable Star Periods from Unevenly Sampled Light Curve Data

Daniel Lichtblau*
Wolfram Research
Champaign
USA
danl@wolfram.com

Abstract

A problem of interest in astronomy is determining the period of variable stars. Data collection is of necessity irregular (can only sample on clear nights) and noisy (from light pollution, atmospheric differences, etc.) We describe several ways in which period estimation can be performed on such data. Some are by now classical (from 60's-80's). One newer method will use Diophantine approximation.

*Jeffrey Bryant

Structured low-rank approximation: Theory, algorithms, and applications

Ivan Markovsky
Department ELEC
Vrije Universiteit Brussel
Belgium
imarkovs@vub.ac.be

Abstract

Mathematical engineering continuously addresses new applications and solves new problems. The expansion of existing methods and applications makes it difficult to maintain a common theoretical framework. This talk shows the potential of the structured low-rank approximation setting to unify problems of data modeling from diverse application areas. An example treated in more details in the presentation is identification of a linear time-invariant system from observed trajectories of the system. We present an optimization method based on the variable projection principle. The method can deal with data with exact and missing (unknown) values. Problems with exact and missing values occur in data driven simulation and control—a new trend of model-free methods for system dynamical analysis and control.

Reference: <http://slra.github.io/>

Well conditioned rational functions approximants versus numerically co-prime polynomials

Ana C. Matos*

Laboratoire Painlevé UMR 8524, UFR Mathématiques

Université de Lille, France

{bbecker,matos}@math.univ-lille1.fr

Abstract

Rational functions like for instance Padé approximants play an important role in signal processing, sparse interpolation and exponential analysis. However, for a successful modeling with help of rational functions we want to make sure that there is no "similar" rational function being degenerate, i.e., having strictly smaller degree of both degrees of numerator and denominator. In particular, we prefer having rational functions without Froissart doublets (i.e., roots close to a pole), and without spurious poles (i.e., simple poles having small residuals).

In a recent paper [1] we showed that, provided that the Sylvester matrix built with the coefficients of the numerator and denominator is well-conditioned, the corresponding rational function has neither Froissart doublets nor spurious poles, and this is also true to sufficiently "close" rational functions. Here closeness is measured with two different metrics, in terms of the chordal distance of the values on the unit disk, or in terms of the distance of normalized coefficient vectors. The paper [1] also contained a comparison of these two metrics.

In [2] the authors introduced a measure for numerical coprimeness representing the minimal distance in the coefficient vector metric to a couple of degenerate polynomials (with a joint root allowing for canceling the fraction). They also showed that if the underlying Sylvester matrix is well-conditioned then a couple of polynomials is numerically coprime, the reciprocal being wrong.

The aim of this talk is to provide precise inequalities implying that also the larger class of rational functions with numerator and denominator being numerically coprime do not have neither Froissart doublets nor spurious poles.

This is a joint work with Bernd Beckermann and George Labahn.

References

- [1] B. Beckermann, A. Matos, Algebraic properties of robust Padé approximants *Journal of Approx. Theory* **190**, 91-115 (2015)
- [2] B. Beckermann, G. Labahn, When are two numerical polynomials relatively prime? *J. Symbolic Computations* **26**, 677-689 (1998).

*Supported in part by the Labex CEMPI (ANR-11-LABX-0007-01).

Using noise to detect faint signals: tricks with Padé approximants to Z-transforms

Luca Perotti* Daniel Bessis[†] Daniel Vrinceanu

Texas Southern University

Houston, TX

USA

{perottl, bessisd, vrinceanud}@tsu.edu

Tania Regimbau

Observatoire de la Côte d'Azur

Nice

France

tania.regimbau@oca.eu

Abstract

Adding small amounts of noise is a recognized method to stabilize Padé approximants; due to the nonlinearity of the Padé approximant, larger amounts of noise can be added to generate different time series and thus increase the statistics when detection probabilities are low. Recently we proposed a new technique based on the observation that the presence of even a weak signal significantly perturbs the universal properties of noise poles and zeros of the Padé approximants to the Z-transform of a data series. For data from two channels, combined in a single complex sequence, the different behavior of poles corresponding to complex noise and poles corresponding to coherent signal can also be used as a signature of the presence of a signal in heavy noise.

*Thanks to my co-workers

[†]Sponsored by NASA

The generalized Prony method and its application I and II

Thomas Peter

Institute for Mathematics, University of Osnabrück, Germany
thomas.peter@uni-osnabrueck.de

Gerlind Plonka

Institute for Numerical and Applied Mathematics
University of Göttingen, Germany
plonka@math.uni-goettingen.de

Abstract

In this paper, we want to present a new very general approach for the reconstruction of sparse expansions of eigenfunctions of suitable linear operators. This approach provides us with a tool to unify all Prony-like methods on the one hand and to essentially generalize the Prony approach on the other hand. Thus it will establish a much broader field of applications of the method. In particular, we will show that all well-known Prony-like reconstruction methods for exponentials and polynomials known so far, can be seen as special cases of this approach. For example, the new insight into Prony-like methods enables us to derive new reconstruction algorithms for orthogonal polynomial expansions including Jacobi, Laguerre, and Hermite polynomials. The approach also applies to finite dimensional vector spaces, and we derive a deterministic reconstruction method for M -sparse vectors from only $2M$ measurements.

The talk will be split into two parts given by the two authors. In the **first part** we concentrate on deriving the new general approach to apply Prony's method to sparse expansions of eigenfunctions of linear operators and present the close connection to the well-known Prony-method.

The **second part** of the talk is especially dedicated to the advantages that the new more general insights give us for applications, as e.g. the use of different linear operators, the influence of the choice of functionals in case of noisy data and further numerical issues.

Efficient spectral estimations by MUSIC and related algorithms

Daniel Potts* Toni Volkmer

Faculty of Mathematics, Technische Universität Chemnitz

D–09107 Chemnitz, Germany

{potts, toni.volkmer}@mathematik.tu-chemnitz.de

Manfred Tasche

Institute of Mathematics, University of Rostock

D–18051 Rostock, Germany

manfred.tasche@uni-rostock.de

Abstract

In the spectral estimation, one has to determine all parameters of a univariate resp. multivariate exponential sum h , if only finitely many (noisy) sampled data of h are given. A frequently used method for spectral estimation is the known MUSIC algorithm. Another popular methods are ESPRIT and the approximate Prony method (APM). We show that both MUSIC and APM are based on an orthogonal projection onto a so-called noise space, whereas ESPRIT uses an orthogonal projection onto the orthogonal complement of the noise space, the so-called signal space. These orthogonal projections can be constructed by (partial) singular value decomposition or QR decomposition of a rectangular Hankel matrix formed by the given sampled data of h .

In this talk, we describe that MUSIC and the related algorithms can be efficiently realized by sampling of h on special grids and using sparse fast Fourier transforms. Numerical experiments illustrate the procedure.

*Sponsored by German Research Foundation within the project PO 711/10-2.

Using univariate algorithms to solve multivariate problems

Daniel S. Roche
United States Naval Academy
Annapolis, Maryland, USA
roche@usna.edu

Abstract

A key feature of sparse interpolation algorithms is that their complexity should scale nicely (often linearly) with the number of variables in the unknown function. In fact, such algorithms can usually be decomposed into two parts: a “base case” univariate interpolation algorithm, and a method to reduce a given multivariate problem to one or more instances of a univariate one.

We will look at both historical and very recent approaches to the second part, the multivariate-to-univariate reduction. As has been frequently observed, many of these reductions are essentially orthogonal to the choice of underlying univariate algorithm, allowing for a wide range of hybrid approaches — not all of which are equally effective. We will examine the strengths and weaknesses of the various variable reduction strategies, and aim to give some insights into how they may be most effectively chosen and applied to new problems.

New approaches to Vector-Valued Rational Interpolation

Avram Sidi

Computer Science Department

Technion - Israel Institute of Technology

Haifa 32000, Israel

asidi@cs.technion.ac.il

<http://www.cs.technion.ac.il/~asidi>

Abstract

We discuss some recent vector-valued rational interpolation procedures for vector-valued functions $F(z)$, $F : \mathbb{C} \rightarrow \mathbb{C}^N$. The interpolants produced by these procedures are all of the simple form

$$R_{p,k}(z) = \frac{U_{p,k}(z)}{V_{p,k}(z)} = \frac{\sum_{j=0}^k c_j \psi_{1,j}(z) G_{j+1,p}(z)}{\sum_{j=0}^k c_j \psi_{1,j}(z)}.$$

Here

$$\psi_{m,n}(z) = \prod_{r=m}^n (z - \xi_r), \quad n \geq m \geq 1; \quad \psi_{m,m-1}(z) = 1, \quad m \geq 1,$$

and $G_{m,n}(z)$ is the vector-valued polynomial of interpolation to $F(z)$ at the points ξ_i , $m \leq i \leq n$. The c_j are scalars, and they are determined in different ways by the different methods. As such, $R_{p,k}(z)$ interpolates $F(z)$ at ξ_i , $1 \leq i \leq p$, in the generalized Hermite sense.

We first discuss the algebraic properties of these interpolants, namely, their uniqueness, symmetry, and reproducing properties. We next discuss their use in approximating vector-valued meromorphic functions $F(z)$ in the complex plane.

Next, choosing the interpolation points appropriately, for $p \rightarrow \infty$ and k fixed, we derive de Montessus type convergence results for the interpolants and Koenig type convergence results for their poles and residues, which show that these interpolants, despite their simple appearance, are effective approximation tools. Especially interesting Koenig type results are obtained when the residues of $F(z)$ form a mutually orthogonal set. (Note that, for any type of rational interpolation problem, whether scalar or vector, the crucial test for deciding whether these are useful approximation tools is the existence of de Montessus and Koenig type theories.)

Finally, we consider the fully confluent case in which all interpolation points ξ_i coincide, and show the connection of the resulting interpolants with Krylov subspace methods.

References

- [1] A. Sidi. Rational approximations from power series of vector-valued meromorphic functions. *J. Approx. Theory*, 77:89–111, 1994.

- [2] A. Sidi. Application of vector-valued rational approximation to the matrix eigenvalue problem and connections with Krylov subspace methods. *SIAM J. Matrix Anal. Appl.*, 16:1341–1369, 1995.
- [3] A. Sidi. A new approach to vector-valued rational interpolation. *J. Approx. Theory*, 130:177–187, 2004.
- [4] A. Sidi. Algebraic properties of some new vector-valued rational interpolants. *J. Approx. Theory*, 141:142–161, 2006.
- [5] A. Sidi. A de Montessus type convergence study for a vector-valued rational interpolation procedure. *Israel J. Math.*, 163:189–215, 2008.
- [6] A. Sidi. A de Montessus type convergence study of a least-squares vector-valued rational interpolation procedure. *J. Approx. Theory*, 155:75–96, 2008.
- [7] A. Sidi. A de Montessus type convergence study of a least-squares vector-valued rational interpolation procedure II. *Comput. Methods Funct. Theory*, 10:223–247, 2010.

Towards simplified construction of subresultant matrix of multiple univariate polynomials

Akira Terui
University of Tsukuba
Tsukuba 305-8571
Japan
terui@math.tsukuba.ac.jp

Abstract

For three or more inputs of univariate polynomials with the real coefficients, we discuss a new construction of subresultant-like matrix which enable us to estimate the degree of the greatest common divisor (GCD) of the input polynomials from its rank. Such matrix is used in approximate GCD algorithms using optimization techniques with its degree is given in advance, especially for constructing constraints. Therefore, in these algorithms, it is important to construct the matrix in a more simplified form to make the overall algorithm more efficient. In this talk, for those purposes, we discuss towards a proposal of a new simplified construction of the matrix.

Hankel and Quasi-Hankel low-rank matrix completion: a convex relaxation

Konstantin Usevich Pierre Comon*

CNRS, GIPSA-Lab,

F-38000 Grenoble,

France

firstname.lastname@gipsa-lab.fr

Abstract

The completion of matrices with missing values under the rank constraint is a non-convex optimization problem. A popular convex relaxation is based on minimization of the nuclear norm (sum of singular values) of the matrix. For this relaxation, an important question is when the two optimization problems lead to the same solution. This question was addressed in the literature mostly in the case of random positions of missing elements and random known elements. In this contribution, we analyze the case of structured matrices with fixed pattern of missing values, in particular, the case of Hankel and quasi-Hankel matrix completion, which appears as a subproblem in the computation of symmetric tensor canonical polyadic decomposition. Similar matrix completion problems appear in other applications, where a function can be approximated as a sum of complex exponentials (time series analysis, medical imaging). We extend existing results on completion of rank-one real Hankel matrices to completion of rank- r complex Hankel and quasi-Hankel matrices.

*This work is supported by ERC AdG-2013-320594 DECODA.

A deterministic sparse FFT algorithm for vectors with short support

Katrin Wannenwetsch
Institute for Numerical and Applied Mathematics
Göttingen University
Germany
k.wannenwetsch@math.uni-goettingen.de

Abstract

It is well known that usual FFT algorithms for the discrete Fourier transform of a vector of length N require $\mathcal{O}(N \log N)$ arithmetical operations. Within the last years, there has been a great interest in sublinear time Fourier algorithms for sparse vectors.

In this talk we consider the special case where a signal $\mathbf{x} \in \mathbb{C}^N$ is known to vanish outside a support interval of length $m < N$. If the support length m of \mathbf{x} or a good bound of it is a-priori known we derive a sublinear algorithm to compute \mathbf{x} from its discrete Fourier transform $\hat{\mathbf{x}} \in \mathbb{C}^N$. The proposed algorithm is deterministic and numerically stable.

In case of exact Fourier measurements we require only $\mathcal{O}(m \log m)$ arithmetical operations. For noisy measurements, we propose a stable $\mathcal{O}(m \log N)$ algorithm.

This is joint work with Gerlind Plonka.

Sparsity with Symbolic Polynomials

Stephen M. Watt
University of Western Ontario
London, Ontario
Canada N6A 5B7
Stephen.Watt@uwo.ca

Abstract

We are interested in algorithms for “symbolic polynomials”, that is multivariate polynomials generalized so the exponents are themselves integer-valued multivariate polynomials, for example $x^{n^2/2-nm/2} - y^m$. These objects may be used to model parameterized families of Laurent polynomials, with integer evaluations of the exponent variables giving specific Laurent polynomials. We have shown elsewhere that when polynomials with coefficients in a particular ring form a unique factorization domain, then so do the corresponding symbolic polynomials. We have given algorithms to compute their GCDs and factorizations in this case. Some of these algorithms rely on reduction to algorithms on sparse polynomials with many more variables, as will be explored in this talk. We additionally describe some new directions on Groebner bases for symbolic polynomials.

Reconstruction of Structured Functions from Sparse Fourier Data

Marius Wischerhoff
Institute for Numerical and Applied Mathematics
Göttingen University
Germany
m.wischerhoff@math.uni-goettingen.de

Abstract

In several scientific areas, such as radio astronomy, computed tomography, and magnetic resonance imaging, the reconstruction of structured functions from the knowledge of samples of their Fourier transform is a common problem. For the analysis of the examined object, it is important to reconstruct the underlying original signal as exactly as possible. We aim to uniquely recover structured functions from only a small number of Fourier samples. For this purpose, the Prony method, which is a deterministic method for the recovery of sparse trigonometric functions, is used as key instrument to derive algorithms for unique recovery by means of a smallest possible set of Fourier data.

We will give an overview of reconstruction results for different function classes, and we will consider two classes in detail.

First, we will examine linear combinations of N non-uniform shifts of a given bivariate function. Here, the unknown shift parameters and corresponding coefficients in the linear combination are recovered from sparse Fourier data. Unique recovery of the parameters is possible by using only $3N + 1$ Fourier samples on three lines through the origin. For this purpose, two predetermined lines are considered, while the third sampling line is chosen dependently on the results obtained by employing the samples from the first two lines. The presented approach can be generalized to the case of d -variate functions with $d > 2$.

Secondly, we turn to the reconstruction of polygonal shapes in the real plane. Here, a convex or non-convex polygonal domain D with N vertices is considered. It is shown that the vertices and their order can be reconstructed by taking $3N$ samples of the Fourier transform of the characteristic function of the polygonal domain D . Again, two predetermined sampling lines and an appropriately chosen third line are considered.

Semidefinite Representations of Noncompact Convex Sets

Lihong Zhi*

Key Laboratory of Mathematics Mechanization
Academy of Mathematics and System Sciences
CAS, Beijing 100190, China
lzhi@mmrc.iss.ac.cn

Abstract

We consider the problem of the semidefinite representation of a class of noncompact basic semialgebraic sets. We introduce the conditions of pointedness and closedness at infinity of a semialgebraic set and show that under these conditions our modified hierarchies of nested theta bodies and Lasserre's relaxations converge to the closure of the convex hull of S . Moreover, if the PP-BDR property is satisfied, our theta body and Lasserre's relaxation are exact when the order is large enough; if the PP-BDR property does not hold, our hierarchies converge uniformly to the closure of the convex hull of S restricted to every fixed ball centered at the origin. We illustrate through a set of examples that the conditions of pointedness and closedness are essential to ensure the convergence. Finally, we provide some strategies to deal with cases where the conditions of pointedness and closedness are violated.

*Thanks to my co-authors: Feng Guo and Chu Wang. Lihong Zhi is supported by NKBRPC 2011CB302400, the Chinese National Natural Science Foundation under grants 91118001.

List of Participants

John Abbott (University of Genova, IT)
Fredrik Andersson (Lund University, SE)
Andrew Arnold (University of Waterloo, CA)
Dmitry Batenkov (Technion – Haifa, IL)
Bernhard Beckermann (University of Lille, FR)
Matteo Biani (University of Antwerp, BE)
Adhemar Bultheel (KU Leuven, BE)
Mathieu Collowald (INRIA Sophia Antipolis – Méditerranée, FR)
Annie Cuyt (University of Antwerp, BE)
Stefano de Marchi (University of Padova, IT)
Pier Luigi Dragotti (Imperial College London, GB)
Jürgen Gerhard (Maplesoft – Waterloo, CA)
Mark Giesbrecht (University of Waterloo, CA)
Karlheinz Gröchenig (Universität Wien, AT)
Lasse Holmström (University of Oulu, FI)
Evelyne Hubert (INRIA Sophia Antipolis – Méditerranée, FR)
Erich Kaltofen (North Carolina State University – Raleigh, US)
Debasis Kundu (Indian Inst. of Technology – Kanpur, IN)
Stefan Kunis (Universität Osnabrück, DE)
George Labahn (University of Waterloo, CA)
Wen-shin Lee (University of Antwerp, BE)
David Levin (Tel Aviv University, IL)
David Li (The University of Strathclyde – Glasgow, GB)
Daniel Lichtblau (Wolfram Research – Champaign, US)
Ivan Markovsky (Free University of Brussels, BE)
Ana C. Matos (Lille I University, FR)
Katharine M. Mullen (UCLA, US)
Luca Perotti (Texas Southern University – Houston, US)
Thomas Peter (Universität Osnabrück, DE)
Gerlind Plonka-Hoch (Universität Göttingen, DE)
Daniel Potts (TU Chemnitz, DE)
Daniel Roche (U.S. Naval Academy – Annapolis, US)
Avraham Sidi (Technion – Haifa, IL)
Manfred Tasche (Universität Rostock, DE)
Akira Terui (University of Tsukuba, JP)
Konstantin Usevich (GRIPSA Lab – Saint Martin d’Hères, FR)
Ulrich von der Ohe (Universität Osnabrück, DE)
Madhavi Wadekar (University of Antwerp, BE)
Katrin Wannenwetsch (Universität Göttingen, DE)
Stephen M. Watt (University of Western Ontario – London, CA)
Marius Wischerhoff (Universität Göttingen, DE)
Yosef Yomdin (Weizmann Institute – Rehovot, IL)
Lihong Zhi (MMRC – Beijing, CN)